

## EXCITATION OF ISOMERIC STATES IN THE REACTIONS $(\gamma, n)$ AND $(n, 2n)$ ON THE $^{81}\text{Br}$ NUCLEUS

*S. R. Palvanov<sup>a</sup>, F. R. Egamova<sup>a</sup>, A. Kh. Ramazanov<sup>a</sup>,  
G. S. Palvanova<sup>a</sup>, A. Kh. Inoyatov<sup>b, c</sup>*

<sup>a</sup> National University of Uzbekistan, Tashkent

<sup>b</sup> Joint Institute for Nuclear Research, Dubna

<sup>c</sup> Institute of Nuclear Physics of AS RUz, Tashkent

The cross sections for the formation of isomeric states in the reactions  $(\gamma, n)$  and  $(n, 2n)$  on the  $^{81}\text{Br}$  nucleus were measured by the induced activity method. The energy dependence of the isomeric ratio of the reaction yields  $(\gamma, n)$  in the energy range 14–35 MeV with a step of 1 MeV was obtained. The experimental results are compared with the data of other works and the TALYS-1.6 calculation.

Методом наведенной активности измерены сечения образования изомерных состояний в реакциях  $(\gamma, n)$  и  $(n, 2n)$  на ядре  $^{81}\text{Br}$ . Получены энергетические зависимости изомерного отношения выходов реакций  $(\gamma, n)$  в области энергий 14–35 МэВ с шагом 1 МэВ. Результаты экспериментов сравниваются с данными других работ и расчетом TALYS-1.6.

PACS: 24.30.Cz; 25.20.Dc

Received on May 31, 2021.